



Reply to comment by Y. Rolland et al. on "Alpine thermal and structural evolution of the highest external crystalline massif: The Mont Blanc"

P.H. Leloup, N Arnaud, R Lacassin, E.R Sobel

► To cite this version:

P.H. Leloup, N Arnaud, R Lacassin, E.R Sobel. Reply to comment by Y. Rolland et al. on "Alpine thermal and structural evolution of the highest external crystalline massif: The Mont Blanc". *Tectonics*, 2007, 26 (2), pp.TC2016. 10.1029/2006TC002022 . insu-01285574

HAL Id: insu-01285574

<https://hal-insu.archives-ouvertes.fr/insu-01285574>

Submitted on 10 Mar 2016

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Reply to comment by Y. Rolland et al. on “Alpine thermal and structural evolution of the highest external crystalline massif: The Mont Blanc”

P. H. Leloup,¹ N. Arnaud,² R. Lacassin,³ and E. R. Sobel⁴

Received 20 July 2006; revised 19 January 2007; accepted 29 January 2007; published 26 April 2007.

Citation: Leloup, P. H., N. Arnaud, R. Lacassin, and E. R. Sobel (2007), Reply to comment by Y. Rolland et al. on “Alpine thermal and structural evolution of the highest external crystalline massif: The Mont Blanc,” *Tectonics*, 26, TC2016, doi:10.1029/2006TC002022.

1. Introduction

[1] *Leloup et al.* [2005] discussed the Cenozoic structural evolution of the Mont Blanc and Aiguilles Rouges ranges by combining new structural, $^{40}\text{Ar}/^{39}\text{Ar}$, and fission track data with published P-T estimates and geochronological data. Our main conclusions were (1) Alpine exhumation of the Aiguilles Rouges was limited to the thickness of the overlying nappes (~ 10 km), while rocks now outcropping in the Mont Blanc have been exhumed 15 to 20 km. (2) Uplift of the two massifs started ~ 22 Myr ago; while at 12 Ma, the Mont Blanc shear zone (MBSz), a reverse fault with a slight right-lateral component, initiated bringing the Mont Blanc above the Chamonix synclinorium and the Aiguilles Rouges; total vertical throw on the MBSz is between 4 and 8 km. (3) Fission track data suggest that relative motion between the Aiguilles Rouges and the Mont Blanc stopped 4 Myr ago. Since that time, uplift of the Mont Blanc has mostly taken place along the Mont Blanc back thrust, a steep north dipping fault zone bounding the southern flank of the range. (4) The highest summits are located where the back thrust intersects the MBSz. (5) Exhumation of the Mont Blanc and Aiguilles Rouges occurred toward the end of motion on the Helvetic basal décollement (HBD) at the base of the Helvetic nappes. Uplift is linked with a deeper, more external thrust that induced the formation of the Jura arc.

[2] While acknowledging that our paper is “a good step forward in the tectonic comprehension of the Mont Blanc area and provides a good synthesis of preexisting data,”

Rolland et al. [2007] claim that the timing we propose for the thrust and back thrust events is not in agreement with new $^{40}\text{Ar}/^{39}\text{Ar}$ data that they publish in their comment. In fact, they raise two main arguments with our observations/interpretations:

[3] 1. Alpine deformation is penetrative within the Mont Blanc granite and is not accommodated by the two localized shear zones we describe (the SE dipping Mont Blanc shear zone, or MBSz, in the north and the NW dipping back thrust in the south, Figure 1), but by numerous anastomosed shear zones in the way described by *Choukroune and Gapais* [1983] in the Aar massif and *Gourlay* [1986] in the Mont Blanc. All deformations within the Mont Blanc are thus coeval and the Mont Blanc is a transpressive pop-up structure at the rim of a large transpressive fault that runs from the Rhone dextral fault system.

[4] 2. The timing of deformation cannot be obtained through $^{40}\text{Ar}/^{39}\text{Ar}$ thermochronology due to excess argon and intense fluid circulation. They instead provide a minimum age of 16 Ma for the initiation of top to the SE motions on the SE side of the Mont Blanc (back thrust) based on five phengites $^{40}\text{Ar}/^{39}\text{Ar}$ ages from three shear zones (their Figure 3).

[5] We will take the opportunity of this reply to address these two points and, in a third point, we briefly discuss possible deformation models of the Mont Blanc range.

2. Alpine Structures in the Mont Blanc Range

2.1. Are All Structures of the Mont Blanc Granite due to Alpine Deformation?

[6] Some authors ascribe most deformation of the Mont Blanc range to the Variscan orogeny and restrict Alpine deformation to small-scale brittle faults and the Faille d'Angle fault bounding the Mont Blanc granite to the NE (Figure 1a) [e.g., *Bellière*, 1988]. One important point of our work was to emphasize the importance of alpine deformation and shear zones within and at the margins of the range. *Rolland et al.* [2007] go even further, stating that all foliations within the Mont Blanc granite result from Alpine flattening. However, there are indeed Variscan deformation in the Mont Blanc range that must be distinguished from Alpine deformation, the tricky point being that both schistositys have nearly the same directions.

[7] On the NW flank of the Mont Blanc, migmatitic gneisses are crosscut by the Mont Blanc granite and by

¹Laboratoire des Sciences de la Terre, UMR CNRS 5570, Ecole Normale Supérieure, Université Lyon 1 and Ecole Normale Supérieure de Lyon, Lyon, France.

²Laboratoire de Dynamique de la Lithosphère, UMR CNRS 5573, ISTEEM-USTL, Montpellier, France.

³Tectonique et Mécanique de la Lithosphère, Institut de Physique du Globe de Paris, and Université Paris 7, CNRS, Paris, France.

⁴Institut für Geowissenschaften, Universität Potsdam, Potsdam, Germany.




Figure 1

aplitic veins, prior to having been affected by greenschist Alpine deformation [see *Leloup et al.*, 2005, Figures 3b and 3h]. In the absence of radiometric data from the dikes, we assumed that as in many places in the External Crystalline Massifs, they are related with late stages of the granite emplacement and probably have an upper Stephanian age as it is the case in the Pelvoux [e.g., *Strzeczynski et al.*, 2005]. This implies that the migmatitic gneisses are Variscan in age. The geometry of the Variscan deformation is clear in the Aiguilles Rouges, where alpine metamorphism and deformation were mild, and in large boudins preserved from alpine deformation within the MBsz. In these zones, foliations (S1) are steep, strike N335 to N55 (N20 on average) and when present, the stretching lineation plunges relatively shallowly [*Leloup et al.*, 2005, Figure 4b]. Variscan (S1) and alpine (S2) foliations cannot be distinguished simply by their strikes because they are close to each other and both vary significantly around a mean value (compare Figures 4a and 4d of *Leloup et al.* [2005]). We ascribed an alpine age to all greenschist foliations bearing a downdip lineation and a Variscan age to higher temperature foliations without, or with shallowly plunging lineation. Aplitic dikes systematically cut the Variscan foliations, but are affected by the alpine ones.

[8] Within the Mont Blanc granite, a high-temperature foliation (S1) that we termed “magmatic foliation” is defined by the alignment of feldspar porphyroclasts and the flattening of restitic enclaves (Figure 2). S1 strikes N-S to NE-SW, does not bear any lineation and is crosscut by undeformed aplitic dikes (Figure 2). This implies that the penetrative deformation of the Mont Blanc granite (S1) is late Variscan. However, both the granite and the aplitic veins are affected by numerous greenschist shear zones and faults that postdate the magmatic foliation and that most probably corresponds to Alpine deformation.

2.2. Geometry of the Alpine Shear Zones Within the Mont Blanc Granite: One or Several Deformation Phases?

[9] *Rolland et al.* [2007] argue that the geometry of the alpine shear zones in the Mont Blanc is analogous to that of the Aar range described by *Choukroune and Gapais* [1983] and reflects a single phase of NW-SE shortening. Despite several studies [*Bellière*, 1956, 1988; *Bertini et al.*, 1985; *Gourlay*, 1986; *Rolland et al.*, 2003; *Rossi*, 2005], the detailed geometry and the amount of shortening of the deformations affecting the core of Mont Blanc granite is still unclear. This is partly due to access difficulties in the highest part of the range, and partly because structures ranging from mylonites to small brittle faults have been mixed together and inappropriately analyzed with the same tools. For example, *Bertini et al.* [1985] present an analysis

of “striated fault planes” that obviously include numerous ductile shear zones, while *Rossi* [2005, Figure II-3] mixes faults and shear zones to calculate stress directions using *Angelier’s* [1990] method, although this should be restricted to microfaults with minor offsets. In the same way, *Rolland et al.* [2007] identify the stretching lineation as the σ_3 stress axis, an assumption that is clearly incorrect in shear zones [e.g., *Ramsay and Huber*, 1983].

[10] What is clear, however, is that the main alpine shear zones are steep, show reverse sense and roughly parallel the main boundaries of the Mont Blanc range: the \sim N35°E, SE dipping MBsz to the NW and the \sim N50°E NW dipping back thrust to the SE (Figure 1a). At the scale of the range, the shear zones have an upward diverging fan-like geometry on NW-SE cross sections (Figure 1b) [also see *Bertini et al.*, 1985, Figure 3; *Rossi*, 2005, Figure II-4]. Near the MBsz, all alpine shear zones dip SE, implying a top to the NW thrusting of the Mont Blanc [*Gourlay*, 1986; *Leloup et al.*, 2005]. Symmetrically, most shear zones dip to the NW near the back thrust [e.g., *Guermani and Pennacchioni*, 1998]. Such a geometry is fundamentally different to that of the core of the Aar range, where shear zones are anastomosed at the scale of few tens of meters, with conjugate shear zones of opposite dips merging around less deformed lenses [*Choukroune and Gapais*, 1983].

[11] What is important for evaluating the structural history of the Mont Blanc range is to determine whether all shear zones and faults are coeval and result from a single deformation episode. This hypothesis appears to be sustained by the large-scale geometry, with the MBsz and the back thrust resembling conjugate faults. However, these faults are not strictly parallel and, cartographically, the back thrust is much shorter than the MBsz (Figure 1). In the absence of unambiguous relative or absolute timing for the shear zones, several lines of evidence lead us to infer that the back thrust was (or is still) active more recently than the MBsz: (1) the back thrust borders the highest summits of the range (Figure 1), suggesting that it has a strong influence on the topography; (2) the main back thrust contact of the Mont Blanc granite on top of the sedimentary series is brittle, exhibiting cataclastic granite [see *Leloup et al.*, 2005, Figure 3e], thus suggesting that deformation occurred late in the uplift history of the range; (3) at a given altitude (i.e., along the Mont Blanc tunnel) fission track ages get younger toward the SE and reach 2.8 ± 0.5 Ma near the back thrust [see *Leloup et al.*, 2005, Figure 11b]; we took this as evidence for very recent uplift on the back thrust; and (4) in the north of the range, where there is no back thrust, the Triassic unconformity, still visible on the top of the Mont Blanc granite (Figure 1), dips \sim 65° to the east as does the overlying series. This passive tilting of the eastern flank of the Mont Blanc range indicates that, in the northern part

Figure 1. Structure of the Mont Blanc massif. Modified from Figure 2 of *Leloup et al.* [2005]. (a) Structural map of the Mont Blanc massif. The black frame corresponds to Figure 1 of *Rolland et al.* [2007]. Black stars are samples from *Rolland et al.* [2007], MB140, MB94, and MB30 from north to south. White stars are samples from *Kirschner et al.* [1996] 93-29A and 93-29J, and *Crespo-Blanc et al.* [1995], 4. Black circles with gray filling are argon samples from *Leloup et al.* [2005]. (b) Synthetic cross section of the Mont Blanc massif. Note that this section is compatible with recent gravity data of the area [*Masson et al.*, 2002]. The geometry of the shear zones within the granite is crudely depicted.




Figure 2. Late Variscan high-temperature foliation in the Mont Blanc granite. (a) Sketch of an aplitic dike (A) crosscutting the Mont Blanc granite “magmatic” foliation S1. S1 and the dike are crosscut by two alpine shear zones (S2). Eperon des géographes above the Leshaux glacier. Horizontal plane, view from above. Hammer gives scale. (b) Undeformed aplitic dike cutting across a flattened (S1) enclave of the Mont Blanc granite. North bank of the Leshaux glacier under the Talèfre glacier. View toward the east of two perpendicular planes: N20 vertical and N110 30S.

of the range, thrusting that drove vertical movement only occurred along the MBsz. Our assertion was that, more to the south, the geometry was similar prior to the initiation of the back thrust. We thus assumed a late activation of the back thrust with respect to the MBsz on our sequential history, depicted in Figure 12 of *Leloup et al.* [2005]. However, from structural arguments, it cannot be definitively excluded that the MBsz and the Back thrust initiated coevally and that the back thrust remained active until more recently.

3. Timing of Deformations: Thermochronology Versus $^{39}\text{Ar}/^{40}\text{Ar}$ Dating of White Micas

[12] From a compilation of geochronological ages we proposed a global thermochronological history with an onset of significant cooling around 22 Ma [*Leloup et al.*, 2005, Figure 8] that we interpret as the onset of exhumation of the Mont Blanc and Aiguilles Rouges. In the same way, we interpret the high number of LT Kf ages around 12 Ma as representing the timing of initiation of the MBsz. It is true that low-temperature geochronology is difficult in the Mont Blanc because of the strong Variscan inheritance, the relatively mild alpine metamorphism ($\sim 400^\circ\text{C}$) [*Poty et al.*, 1974; *Marshall et al.*, 1998; *Rolland et al.*, 2003], and the fluid circulation overprint. *Rolland et al.* [2007] propose to date the top to the SE deformation by using $^{39}\text{Ar}/^{40}\text{Ar}$ dating of synkinematic white micas. The great advantage of that method is to

associate an age with a given structure, not with a cooling history. However, each approach has its own pitfalls and one must be cautious prior to reaching definite conclusions.

[13] 1. Dated white micas are claimed to have crystallized in pressure shadows of Kf porphyroclasts. This is obviously not the case in the picture provided by *Rolland et al.* [2005, Figure 2], where white micas follow the foliation. This does not prove that these micas did not crystallize during top to the east shearing but suggests that they could have formed during a previous deformation stage and were later reoriented, thus giving the age of another deformation event.

[14] 2. Analyzing aggregates after crushing, despite careful selection, leaves open the possibility of sampling parts of older micas, as it is the case when working on populations. Only direct dating on thin section could avoid this problem. A simple calculation shows that contamination from only 3% of Variscan (~ 300 Ma) white mica with similar K content to the alpine ones will shift a ~ 5 Ma age to ~ 15 Ma. We note that no real plateau was achieved from the phengites and that all samples yield low-temperature ages around 12 Ma, which is attributed to possible Ar loss during later deformation or fluid percolation. However, an intimate mix of ~ 12 Ma old micas with older preserved ones would probably produce the same shape of spectra.

[15] 3. Finally, cooling ages and dating of synkinematic minerals need not be in opposition but should be discussed in concert. Indeed, *Rolland et al.* [2007] agree that the ages of their micas provide a minimum constraint on the age of deformation and possibly record cooling rather than defor-

mation, and their ages (~ 16 Ma) fall at the time we infer for major cooling linked to exhumation. It is possible, therefore, that dated white micas formed during an earlier deformation event, and recorded cooling at ~ 15 Ma associated with a major phase of Mont Blanc exhumation.

[16] In order to validate their ages, *Rolland et al.* [2007] stress that these are close to those obtained on white micas within shear zones of the Mont Blanc sedimentary cover [*Crespo-Blanc et al.*, 1995; *Kirschner et al.*, 1996]. However, the age of 15.5 Ma of *Crespo-Blanc et al.* [1995] corresponds to the end of west verging thrusting of Helvetic sediments on top of the Mont Blanc along the Val Ferret thrust (Figure 1). That thrust has a different strike and an opposite vergence than the Mont Blanc back thrust. In the same way, *Kirschner et al.* [1996] date the end of motion at the base of the NW verging Morcles and Doldenhorn Helvetic nappes (Figure 1). This has no direct connection with the age of initiation of motion along the SE verging Mont Blanc back thrust.

[17] The ~ 16 Ma ages of *Rolland et al.* [2007] could correspond to cooling below $390 \pm 45^\circ\text{C}$ during the initial stage of the alpine Mont Blanc uplift rather than the activation of the Mont Blanc back thrust. Alternatively, considering an initiation of motion on the back thrust at ~ 16 Ma would only imply that exhumation on top of the basal décollement has been accommodated in part by the back thrust since the beginning, not postponed until ~ 4 Ma, as we originally proposed.

4. Alpine Deformations of the Mont Blanc Range: Push-up Within a Dextral Strike-Slip Shear Zone or Culmination Above an Alpine Basal Thrust?

[18] Finally, the last issue invoked by *Rolland et al.* [2007] is to state, without much discussion, that the new ages they present imply that the Mont Blanc is a push-up along a major dextral shear zone, dismissing our interpretation of a culmination above a basal thrust. Note first that the northern half of the range and its southern extremity do not have a symmetrical shape. It is only where the back thrust is present that the Mont Blanc range forms a crustal-scale pop-up (Figure 1). Furthermore, evidence for significant long-lasting (>16 Myr) strike-slip shear in the Mont Blanc range and along its sides are very weak. In any case,

there are no new arguments in the work of *Rolland et al.* [2007] to support major NE-SW dextral strike-slip motion absorbed within the Mont Blanc. Kinematics of the MBsz, with pitches of lineations larger than 70° , suggests a maximum of 2 km of dextral motion compared with the ~ 26 km of NW-SE shortening absorbed in the Jura arc during that time interval [*Leloup et al.*, 2005, and references therein]. Similarly, structural evidence for major right-lateral shear in the SE part of the range is weak. There, the lack of stretching lineations render the precise direction of shearing difficult to constrain, but most shear zones display shear criteria suggesting top to the SE (dip slip) thrusting [*Guermani and Pennacchioni*, 1998]. The brittle faults that affect the granite are mostly parallel to the Mont Blanc back thrust and often show dip slip motions; only a few brittle faults that reactivate schistosity planes in the Val Veni sedimentary series show dextral motion [see *Leloup et al.*, 2005, Figure 6].

[19] The idea of major continuous dextral motion in the Mont Blanc since the middle Miocene still await convincing structural evidences, although a complex connection of the Mont Blanc thrusts with the Rhône-Valais strike-slip zone and with the Simplon normal fault in a way comparable to that described by *Lacassin* [1986] probably exists.

5. Conclusion

[20] To answer to the main points raised by *Rolland et al.* [2007], we make the following conclusions:

[21] 1. Some of the foliations observed in the Mont Blanc granites are pre-Alpine and cannot be used as evidence for alpine penetrative deformation.

[22] 2. NW and SE dipping shear zones in the Mont Blanc granite may be, but are not necessarily, synchronous. They could as well reflect several stages of deformation that may have overlapped in time.

[23] 3. The new $^{39}\text{Ar}/^{40}\text{Ar}$ ages of ~ 16 Ma obtained on white micas are compatible with the global thermal history that we proposed for the Mont Blanc. They might date initiation of the south verging shear zones but would require a more complete geochronologic and structural analysis in order to be fully validated.

[24] 4. The new data do not bring any new arguments in favor of a right-lateral transpression context for the Mont Blanc range since the middle Miocene.

References

- Angelier, J. (1990), Inversion of field data in fault tectonics to obtain the regional stress. III: A new rapid inversion method by analytical means, *Geophys. J. Int.*, 103, 363–376.
- Bellièvre, J. (1956), Caractère de la déformation Alpine dans les schistes cristallins du massif du Mont-Blanc, *Bull. Soc. Geol. Fr.*, 6, 691–698.
- Bellièvre, J. (1988), On the age of mylonites within the Mont-Blanc massif, *Geodin. Acta*, 2, 13–16.
- Bertini, G., M. Marcucci, R. Nevini, P. Passerini, and G. Sguazzoni (1985), Patterns of faulting in the Mt Blanc granite, *Tectonophysics*, 11, 65–106.
- Choukroune, P., and D. Gapais (1983), Strain pattern in the Aar granite (central Alps): orthogneiss developed by bulk inhomogeneous flattening, *J. Struct. Geol.*, 5, 1–10.
- Crespo-Blanc, A., H. Masson, Z. Sharp, M. Cosca, and J. Hunziker (1995), A stable and $^{40}\text{Ar}/^{39}\text{Ar}$ isotope study of a major thrust in the Helvetic nappes (Swiss Alps): Evidence for fluid flow and constraints on the nappe kinematics, *Geol. Soc. Am. Bull.*, 107, 1129–1144.
- Gourlay, P. (1986), La déformation du socle et des couvertures delphino-helvétiques dans la région du Mont-Blanc (Alpes occidentales), *Bull. Soc. Geol. Fr., Ser. II*, 8, 159–169.
- Guermani, A., and G. Pennacchioni (1998), Brittle precursors of plastic deformation in a granite: an ex-

- emple from the Mont-Blanc massif (Helvetic, western Alps), *J. Struct. Geol.*, **20**, 135–148.
- Kirschner, L., M. A. Cosca, H. Masson, and J. C. Hunziker (1996), Staircase $^{40}\text{Ar}/^{39}\text{Ar}$ spectra of fine-grained white mica: Timing and duration of deformation and empirical constraints on argon diffusion, *Geology*, **24**, 747–750.
- Lacassin, R. (1986), Plate-scale kinematics and compatibility of crustal shear zones in the Alps, in *Alpine Tectonics*, edited by M. P. Coward et al., *Geol. Soc. Spec. Publ.*, **45**, 339–352.
- Leloup, P. H., N. Audaud, E. R. Sobel, and R. Lacassin (2005), Alpine thermal and structural evolution of the highest external crystalline massif: The Mont Blanc, *Tectonics*, **24**, TC4002, doi:10.1029/2004TC001676.
- Marshall, D., H. R. Pfeifer, J. C. Hunziker, and D. Kirschner (1998), A pressure-temperature-time path for the NE Mont-Blanc massif: Fluid-inclusion, isotopic and thermobarometric evidence, *Eur. J. Mineral.*, **10**, 1227–1240.
- Masson, F., F. Gal, and P. H. Leloup (2002), Une carte gravimétrique haute résolution du massif du Mont-Blanc: Implications structurales, *C. R. Geosci.*, **334**, 1011–1019.
- Poty, B., H. A. Stalder, and A. M. Weisbrod (1974), Fluid inclusions studies in quartz from fissures of western and central Alps, *Schweiz. Mineral. Petrogr. Mitt.*, **54**, 717–752.
- Ramsay, J. G., and M. I. Huber (1983), *The Techniques of Modern Structural Geology: Strain Analyses*, 307 pp., Elsevier, New York.
- Rolland, Y., S. F. Cox, A. M. Boullier, G. Pennacchioni, and N. Mancktelow (2003), Rare Earth and trace element mobility and fractionation in mid-crustal shear zones: insights from the Mont-Blanc Massif (western Alps), *Earth Planet. Sci. Lett.*, **214**, 203–219.
- Rolland, Y., M. Corsini, M. Rossi, S. F. Cox, G. Pennacchioni, N. Mancktelow, and A. M. Boullier (2007), Comment on “Alpine thermal and structural evolution of the highest external crystalline massif: The Mont Blanc” by P. H. Leloup, N. Audaud, E. R. Sobel, and R. Lacassin, *Tectonics*, doi:10.1029/2006TC001956, in press.
- Rossi, M. (2005), Déformation, transferts de matière et de fluide dans la croûte continentale: application aux massifs cristallins externes des Alpes, Ph.D. thesis, 388 pp., Univ. of Grenoble, Grenoble, France.
- Strzeczynski, P., S. Guillot, G. Courrioux, and P. Ledru (2005), Modélisation géométrique 3D des granites stéphaniens du massif du Pelvoux (Alpes, France), *C. R. Geosci.*, **337**, 1284–1292.

N. Arnaud, ISTEEM-USTL, Place Eugène Bataillon, F-34095 Montpellier Cedex 5, France.

R. Lacassin, Institut de Physique du Globe de Paris, 4 place Jussieu, F-75252 Paris Cedex 05, France.

P. H. Leloup, Laboratoire des Sciences de la terre, UMR CNRS 5570, Université Lyon 1, Lyon, F-69003 France. (herve.leloup@univ-lyon1.fr)

E. R. Sobel, Institut für Geowissenschaften, Universität Potsdam, Karl Liebknecht Strasse 24, D-14476 Golm, Germany.